

表-5 昭和52年度二酸化硫黄濃度

(単位: ppb)

測定局	暖房			非暖房			全年			日平均 値2% 除外値	備考
	昼	夜	全日	昼	夜	全日	昼	夜	全日		
利府	2.5	2.3	2.4	4.3	3.1	3.7	3.5	2.8	3.1	10	○
塩釜	4.7	3.9	4.3	6.8	4.3	5.5	5.9	4.1	5.0	15	○
七ヶ浜-1	10.0	4.0	7.0	5.3	3.9	4.6	7.3	3.9	5.6	16	○
多賀城-1	2.6	2.2	2.4	4.6	2.9	3.8	3.8	2.6	3.2	9	○
名取	13.7	11.6	12.6	15.7	13.9	14.8	14.9	13.0	13.9	26	
岩沼	15.6	13.7	14.6	14.8	13.5	14.1	15.1	13.5	14.3	24	
国設仙台	19.7	10.3	15.0	15.5	14.4	14.9	17.2	12.7	14.9	30	
多賀城-2	13.0	11.5	12.3	13.6	11.1	12.3	13.4	11.3	12.3	24	
七ヶ浜-2	9.8	8.3	9.0	11.2	10.6	10.9	10.6	9.6	10.1	15	
柴田	10.1	8.3	9.2	12.6	11.2	11.9	11.6	10.0	10.8	20	
泉	1.8	1.7	1.8	3.7	3.1	3.4	2.9	2.6	2.7	7	○
白石	3.4	3.4	3.4	3.9	3.0	3.5	3.7	3.2	3.5	10	○
矢本-1	8.6	8.3	8.4	11.1	10.6	10.9	10.1	9.6	9.9	17	
石巻-1	10.1	9.4	9.8	12.4	10.7	11.5	11.4	10.2	10.8	19	
古川	13.1	10.8	12.0	13.5	12.0	12.8	13.4	11.5	12.4	22	
矢本-2	7.9	7.7	7.8	10.2	10.1	10.1	9.2	9.1	9.1	19	
松島	3.9	3.2	3.5	3.7	2.9	3.3	3.8	3.0	3.4	8	○
大和	3.7	2.8	3.3	4.1	3.6	3.8	3.9	3.3	3.6	8	○
国設麓岳	3.6	3.5	3.5	2.4	2.4	2.4	2.9	2.8	2.9	8	○
気仙沼	4.2	3.6	3.9	4.8	3.3	4.0	4.5	3.4	4.0	10	○
高砂	11.9	10.2	11.1	19.5	15.6	17.6	15.6	12.8	14.2	29	
岩切	8.5	8.0	8.2	12.0	10.1	11.0	10.4	9.1	9.7	15	
鶴ヶ谷	12.2	10.0	11.1	11.5	9.9	10.7	11.9	9.9	10.9	18	
衛試	21.2	15.2	18.2	18.2	15.0	16.5	19.4	15.1	17.2	28	
東長町	13.0	10.0	11.5	14.4	11.8	13.1	13.6	10.8	12.2	21	
中山	10.2	8.2	9.2	12.2	10.0	11.1	11.1	9.1	10.1	15	
中野	11.1	8.2	9.6	15.0	12.6	13.8	13.0	10.2	11.6	19	
七郷	10.0	8.9	9.4	8.2	7.8	8.0	9.2	8.4	8.8	13	
山田	16.9	8.8	12.8	15.2	10.0	12.6	16.0	9.4	12.7	-	

注 備考欄○印は高感度型測定器による。

測定局	暖房	
	昼	夜
利府	19.3	21.5
塩釜	35.8	31.7
七ヶ浜-1	12.9	11.4
多賀城-1	21.0	23.3
岩沼	26.9	26.1
国設仙台	36.2	33.3
多賀城-2	23.3	20.9
柴田	39.2	43.0
白石	26.8	27.7
矢本-1	19.7	22.4
石巻-1	34.7	37.1
古川	35.5	30.3
松島	19.9	26.8
国設麓岳	4.0	3.3
気仙沼	17.2	14.2
高砂	39.6	47.2
岩切	35.3	36.7
鶴ヶ谷	30.0	37.7
衛試	1.6	26.3
東長町	45.2	47.9
中山	24.2	19.3
中野	25.9	32.0
七郷	46.4	55.2
山田	42.0	0